



BROILER MACRO (medicated)

AMINO ACID MINERAL TRACE MINERAL VITAMIN CONCENTRATE FOR BROILERS
FOR HOME MIXING OF BROILER FEEDS

KEY ADVANTAGES

- Highly concentrated concentrate
- Make maximal use of own/local ingredients
- Contains a growth promoter and a coccidiostat
- Contains phytase enzyme which improves feed conversion

EXAMPLES OF FEEDS

| INGREDIENTS | STARTER | GROWER | FINISHER |
|---------------------------|---------|--------|----------|
| Maize meal | 575 | 625 | 700 |
| Wheat bran | 66 | 66 | 66 |
| Soybean oilcake meal | 325 | 275 | 200 |
| Feed lime | 15 | 15 | 15 |
| Broiler Macro (Medicated) | 19 | 19 | 19 |
| Total (kg) | 1000 | 1000 | 1000 |
| Composition (%) | | | |
| Protein | 21.4 | 19.4 | 16.5 |
| Lysine | 1.25 | 1.11 | 0.91 |
| Methionine | 0.49 | 0.46 | 0.42 |
| Energy (MJ/kg) | 12.1 | 12.3 | 12.6 |
| Calcium | 0.99 | 0.98 | 0.95 |
| Phosphorous | 0.73 | 0.71 | 0.68 |
| Phosphorous (available) | 0.41 | 0.40 | 0.38 |

FEEDING RECOMMENDATIONS

- Mix strictly according to recommendations. Consult a Feedtek nutritionist if alternative ingredients are available
- Feed starter feed for the first 7 to 14 days
- Feed grower feed from day 7 to 14 until about day 28
- Feed finisher feed from about day 28 until slaughter
- Pullets – feed grower feed until 6 weeks of age and finisher feed from 6 to 12 weeks of age
- Contains ionophores - may not be fed to horses or game
- An alternative medicament free product is also available

